




Apomorphine Induced Rotational Response in Parkinsonian Rats Engrafted with KDR Selected Stem Cells




BEST AVAILABLE COPY

FIG. 1



Electron of injected electron beam. MIPM looked
at 1000 ft. in right of the image, and cells
appear to be migrating towards lower end of the



BEST AVAILABLE COPY



FIG. 3





BEST AVAILABLE COPY

FIG. 4

A



B




6/21



BEST AVAILABLE COPY

FIG. 6



BEST AVAILABLE COPY


FIG. 7a



BEST AVAILABLE COPY


FIG. 7b

9/21



BEST AVAILABLE COPY

FIG. 7c




BEST AVAILABLE COPY

FIG. 7d



BEST AVAILABLE COPY

FIG. 7e




BEST AVAILABLE COPY
FIG. 7f



BEST AVAILABLE COPY

FIG. 7g



BEST AVAILABLE COPY

FIG. 7h






FIG. 7i




BEST AVAILABLE COPY

FIG. 7j




BEST AVAILABLE COPY

FIG. 7k



BEST AVAILABLE COPY


FIG. 71



BEST AVAILABLE COPY


FIG. 7m

20/21



BEST AVAILABLE COPY

FIG. 7n



BEST AVAILABLE COPY

FIG. 8